

• • • • • • • • • •
• • • • • • • • • •

Measurements of ϕ_1^{eff} from $K_S K_S K_S$, $K_S \pi^0 \pi^0$ and $K^0 \pi^0$

M. Fujikawa
The Belle Collaboration



Introduction

Δt : proper time difference

Δm : mass difference

η_{CP} : CP eigenvalue

$$\lambda = \frac{q}{p} \frac{A(\bar{B}^0 \rightarrow f)}{A(B^0 \rightarrow f)} \approx \eta_{CP} e^{-i2\phi_1}$$

- CP Asymmetry


$$A(\Delta t) = \frac{\Gamma(\bar{B}^0(\Delta t) \rightarrow f_{CP}) - \Gamma(B^0(\Delta t) \rightarrow f_{CP})}{\Gamma(\bar{B}^0(\Delta t) \rightarrow f_{CP}) + \Gamma(B^0(\Delta t) \rightarrow f_{CP})}$$

$$= \frac{|\lambda|^2 - 1}{1 + |\lambda|^2} \cos(\Delta m \cdot \Delta t) + \frac{2\Im\lambda}{1 + |\lambda|^2} \sin(\Delta m \cdot \Delta t)$$

$\mathcal{A} (= -C) \sim 0$
 Belle BaBar
 Direct CPV
 SM expectation

$\mathcal{S} : -\eta_{CP} \sin 2\phi_1$
 Mixing-induced CPV

Basic Analysis Procedure



e^- : 8.0 GeV
 e^+ : 3.5 GeV

$\Delta z = \Delta t \beta \gamma c \sim 200 \mu m$
 $\beta \gamma = 0.43 / 0.56$
 Belle / BaBar


Basic Analysis Procedure

- B extracted with M_{bc} , ΔE


$$M_{bc} \equiv m_{ES} \equiv \sqrt{E_{beam}^{*2} - p_B^{*2}}, \quad \Delta E \equiv E_B^* - E_{beam}^*$$

- Main Background

- Continuum event [$e^+e^- \rightarrow q\bar{q}$ (q=u,d,s,c)]
- Separate with Likelihood ratio ($L_{s/b}$) from event shape



$e^+e^- \rightarrow Y(4S) \rightarrow B\bar{B}$
(Spherical)





$e^+e^- \rightarrow q\bar{q}$
(Jet-like)

- Signal extraction


- Multi-dimensions (M_{bc} , ΔE , $L_{s/b}$, ...)
- Extended unbinned maximum likelihood fit

Vertex Reconstruction with K_S

- No primary tracks from B vertex
- Extrapolate K_S track to the Interaction Point
- Events are required to have enough SVD hits for vertexing




SVT structure




SVD structure

Vertex Reconstruction with K_S



The validity is confirmed using the $J/\psi K_S$ control sample.


- ⊕ B^0 Lifetime 1.503 ± 0.036 ps
- ⊕ $\sin 2\phi_1 = +0.68 \pm 0.06$



- $\langle \sigma_z \rangle$ resolution similar to normal modes
- Events without the vertex can still be used to measure $\mathcal{A}(-C)$

$B^0 \rightarrow K_S K_S K_S$

- Dominated by $b \rightarrow s \bar{q} q$ penguin decay
 - Theoretically clean (no u quarks in the final state)



- CP even, regardless of any resonant structure
[T. Gershon and M. Hazumi, PLB 596 163 (2004)]

SM expectation \rightarrow

$$\mathcal{S} = -\sin 2\phi_1$$

$$\mathcal{A} = 0$$

$B^0 \rightarrow K_S K_S K_S$ Signal Yield









465 $M\bar{B}\bar{B}$



PRL 98 (2007) 031802

535 $M\bar{B}\bar{B}$





274 ± 20 $K_S K_S K_S$ signal

185 ± 17 $K_S K_S K_S$ signal

$B^0 \rightarrow K_S K_S K_S$ tCPV result



465 MB \bar{B}




$$\mathcal{A} = -\mathcal{C} = +0.16 \pm 0.17 \pm 0.03$$

$$\mathcal{S} = -0.90 \pm \begin{matrix} 0.20 \\ 0.18 \end{matrix} \pm \begin{matrix} 0.04 \\ 0.03 \end{matrix}$$



PRL 98 (2007) 031802


535 MB \bar{B}



$$\mathcal{A} = +0.31 \pm 0.20 \pm 0.07$$

$$\mathcal{S} = -0.30 \pm 0.32 \pm 0.08$$

$B^0 \rightarrow K_S K_S K_S$ Comparison



$$C_{CP} = -\mathcal{A}$$

BaBar $-0.16 \pm 0.17 \pm 0.03$

Belle $-0.31 \pm 0.20 \pm 0.07$

Average -0.23 ± 0.13

$$\sin 2\phi_1^{\text{eff}} = -\mathcal{S}$$


BaBar $0.90 \pm 0.20 \pm 0.04$
 0.18 ± 0.03

Belle $0.30 \pm 0.32 \pm 0.08$

Average 0.74 ± 0.17

$B^0 \rightarrow K_S \pi^0 \pi^0$

- Dominated by $b \rightarrow s \bar{q} q$ penguin decay



- CP even, regardless of any resonant structure
[T. Gershon and M. Hazumi, PLB 596 163 (2004)]

SM expectation

$$\mathcal{S} = -\sin 2\phi_1$$

$$\mathcal{A} = 0$$







$B^0 \rightarrow K_S \pi^0 \pi^0$ Signal Yield



PRD 76 (2007) 071101
227 MB \bar{B}



arXiv.0708.1845
657 MB \bar{B}



117 ± 27 $K_S \pi^0 \pi^0$ signal

307 ± 32 $K_S \pi^0 \pi^0$ signal



$B^0 \rightarrow K_S \pi^0 \pi^0$ tCPV Result



PRD 76 (2007) 071101
227 MB \bar{B}



arXiv.0708.1845
657 MB \bar{B}



LR > 0.9,
good tag


$$\mathcal{A} = -\mathcal{C} = -0.23 \pm 0.52 \pm 0.13$$

$$\mathcal{S} = +0.72 \pm 0.71 \pm 0.08$$

$$\mathcal{A} = -0.17 \pm 0.24 \pm 0.06$$

$$\mathcal{S} = +0.43 \pm 0.49 \pm 0.09$$

$K_S \pi^0 \pi^0$ tCPV Comparison



$$\sin 2\phi_1^{\text{eff}} = -\mathcal{S}$$

BaBar $-0.72 \pm 0.71 \pm 0.08$

Belle $-0.43 \pm 0.49 \pm 0.09$

Average -0.52 ± 0.41

$$C_{CP} = -\mathcal{A}$$

BaBar $0.23 \pm 0.52 \pm 0.13$

Belle $0.17 \pm 0.24 \pm 0.06$

Average 0.18 ± 0.22

$B^0 \rightarrow K^0 \pi^0$

- $A_{CP}(B^0 \rightarrow K^+ \pi^-) \neq A_{CP}(B^+ \rightarrow K^+ \pi^0)$

⊕ ΔA_{CP} puzzle Nature 452, 332-335(2008)

- Isospin sum rule among $B \rightarrow K \pi$ CP asymmetries


M. Gronau, PLB 672(2005)82-88)

$$A_{CP}(K^+ \pi^-) + A_{CP}(K^0 \pi^+) \frac{B(K^0 \pi^+) \tau_0}{B(K^+ \pi^-) \tau_+}$$

$$= A_{CP}(K^+ \pi^0) \frac{2B(K^+ \pi^0) \tau_0}{B(K^+ \pi^-) \tau_+} + A_{CP}(K^0 \pi^0) \frac{B(K^0 \pi^0)}{B(K^+ \pi^-)}$$

- ⊕ Breaking sum rule indicates new physics
- ⊕ Theoretical uncertainty \sim SU(2) breaking



- Both S and A are important



$B^0 \rightarrow K_S \pi^0$ Signal Yield






465 $M\bar{B}$



$556 \pm 32 K_S \pi^0$ signal



657 $M\bar{B}$





$657 \pm 37 K_S \pi^0$ signal


$B^0 \rightarrow K_L \pi^0$ Signal Yield



657 MB \bar{B}



Background subtraction




- First measurement
- M_{bc} calculated from direction of K_L cluster
- $K_L \pi^0$ signal
 285 ± 52 (stat) ± 57 (syst)
 3.7σ (including systematics)

$B^0 \rightarrow K^0 \pi^0$ tCPV result



465 MB \bar{B}




$$\mathcal{A} = -\mathcal{C} = -0.13 \pm 0.13 \pm 0.03$$

$$\mathcal{S} = +0.55 \pm 0.20 \pm 0.03$$



657 MB \bar{B}




$K_S \pi^0 + K_L \pi^0$

$$\mathcal{A} = +0.14 \pm 0.13 \pm 0.06$$

$$\mathcal{S} = +0.67 \pm 0.31 \pm 0.08$$

$B^0 \rightarrow K^0 \pi^0$ Comparison



$$\sin 2\phi_1^{\text{eff}} = \mathcal{S}$$

BaBar $0.55 \pm 0.20 \pm 0.03$

Belle $0.67 \pm 0.31 \pm 0.06$

Average 0.57 ± 0.17

$$C_{CP} = -\mathcal{A}$$


BaBar $0.13 \pm 0.13 \pm 0.03$

Belle $-0.14 \pm 0.13 \pm 0.06$


Average 0.01 ± 0.10

Summary


$K_S K_S K_S C_{CP}$ **HFAG**
CKM2008
PRELIMINARY




$K_S K_S K_S S_{CP}$ **HFAG**
CKM2008
PRELIMINARY




$\pi^0 \pi^0 K_S C_{CP}$ **HFAG**
LP 2007
PRELIMINARY




$\pi^0 \pi^0 K_S S_{CP}$ **HFAG**
LP 2007
PRELIMINARY



$\pi^0 K^0 C_{CP}$ **HFAG**
ICHEP 2008
PRELIMINARY



$\pi^0 K^0 S_{CP}$ **HFAG**
ICHEP 2008
PRELIMINARY



Summary

- Results from Babar and Belle
 - HFAG average shows no significant deviation from SM

	$C_{CP} = -\mathcal{A}$	$\sin 2\phi_1^{\text{eff}}$	
$K_S K_S K_S$	-0.23 ± 0.13	0.74 ± 0.17	← Theoretically clean
$K_S \pi^0 \pi^0$	0.18 ± 0.22	-0.52 ± 0.41	← Anomaly?
$K^0 \pi^0$	0.01 ± 0.10	0.57 ± 0.17	← Sum rule predicts sizable direct CPV

- Super B factory is necessary for these modes
- We need more statistics

Backup


Systematic Errors



	$K_S K_S K_S$		$K_S \pi^0 \pi^0$		$K^0 \pi^0$	
	δS	δA	δS	δA	δS	δA
Vertexing	0.010	0.020	0.011	0.020	0.013	0.022
Flavor tagging	0.012	0.006	0.008	0.005	0.007	0.005
Resolution	0.049	0.016	0.066	0.010	0.063	0.007
Physics	0.001	0.001	0.007	0.001	0.007	0.001
Fit bias	0.024	0.013	0.009	0.004	0.010	0.020
BG fraction	0.057	0.049	0.009	0.001	0.029	0.022
BG dt shape	0.007	0.010	0.046	0.019	0.015	0.006
TSI	0.001	0.042	0.001	0.043	0.014	0.054

Total	0.081	0.071	0.082	0.053	0.06	0.08

KEKB & PEP-II



9 GeV e⁻ x 3.1 GeV e⁺
 Head-on collision

PEP-II (USA)

$\beta\gamma=0.56$

8 GeV e⁻ x 3.5 GeV e⁺
 ±1 mrad crossing

KEKB (Japan)

$\beta\gamma=0.425$

Belle and BaBar Detectors

